

Forward in Health Clinics

February 2008

Patient Diagnoses

| | 2/7/08 | 2/8/08 | 2/9/08 | 2/11/08 | Orphanage | TOTAL | PERCENTAGE* |
|-------------------|--------|--------|--------|---------|-----------|-------|-------------|
| Abdominal pain | 1 | | 1 | 2 | | 4 | 1.0% |
| Abrasions | 1 | 1 | | | | 2 | 0.5% |
| Acne | | 1 | | | | 1 | 0.2% |
| Adenopathy | | 1 | 1 | | | 2 | 0.5% |
| Allergic rhinitis | 2 | | | 1 | | 3 | 0.7% |
| Anemia | 1 | | | | | 1 | 0.2% |
| Asthma | 3 | 1 | 2 | | 1 | 7 | 1.7% |
| Back pain | | | | 1 | | 1 | 0.2% |
| Bronchitis | | 2 | | | | 2 | 0.5% |
| Bronchospasm | 1 | | 1 | 1 | | 3 | 0.7% |
| Chest pain | | | | 1 | | 1 | 0.2% |
| Chicken pox | | 1 | | | | 1 | 0.2% |
| Cold | 8 | 14 | 3 | 1 | | 26 | 6.4% |
| Conjunctivitis | 1 | 1 | 1 | 1 | | 4 | 1.0% |
| Constipation | | 1 | | | | 1 | 0.2% |
| Cough | 3 | 8 | 6 | 3 | | 20 | 4.9% |
| Cradle cap | 1 | | | | | 1 | 0.2% |
| Dandruff | 2 | 4 | 4 | 1 | | 11 | 2.7% |
| Delayed puberty | 1 | | | | | 1 | 0.2% |
| Dental caries | | 3 | | | | 3 | 0.7% |
| Dermatitis | | | | 1 | | 1 | 0.2% |
| Diarrhea | | 1 | 1 | 1 | | 3 | 0.7% |

| | | | | | | | |
|--------------------------|---|---|---|---|---|----|------|
| Dyspepsia | 1 | | | | | 1 | 0.2% |
| Eczema | 7 | 7 | 3 | 5 | | 22 | 5.4% |
| Epistaxis | | 1 | | | | 1 | 0.2% |
| Eye allergy | 1 | | | | | 1 | 0.2% |
| Fever | 2 | 5 | 4 | 8 | | 19 | 4.7% |
| GE Reflux | 1 | 1 | | | | 2 | 0.5% |
| Headaches | 5 | 3 | 2 | | 3 | 13 | 3.2% |
| Hydrocele | | 1 | 1 | | | 2 | 0.5% |
| Hyperdynamic heart | | 1 | | | | 1 | 0.2% |
| Impetigo | | 2 | 1 | 1 | | 4 | 1.0% |
| Knee pain | | | | 1 | | 1 | 0.2% |
| Leg cramps | | | 1 | | | 1 | 0.2% |
| Leg pain | | | | 1 | | 1 | 0.2% |
| Lethargy | 1 | | | | | 1 | 0.2% |
| Lymphadenopathy | | | 1 | | | 1 | 0.2% |
| Malaria | 1 | 2 | | | | 3 | 0.7% |
| Malnutrition | 8 | | | | | 8 | 2.0% |
| Menarche | 1 | 1 | | | | 2 | 0.5% |
| Menstrual cramps | 1 | | 1 | 2 | | 4 | 1.0% |
| Mumps | | | | 1 | | 1 | 0.2% |
| Neck pain | 1 | | | | | 1 | 0.2% |
| Otitis | 1 | 5 | 7 | 4 | | 17 | 4.2% |
| Pica | 1 | | 1 | | | 2 | 0.5% |
| Pneumonia | 1 | | | | | 1 | 0.2% |
| Post chicken pox lesions | 1 | | | | | 1 | 0.2% |
| Pyoderma | 8 | 1 | 1 | 2 | | 12 | 3.0% |
| Rash | 1 | 4 | 4 | 3 | | 12 | 3.0% |
| Scabies | | 7 | 2 | | | 9 | 2.2% |
| Skin Infection | | 6 | 1 | 2 | 1 | 10 | 2.5% |
| Staph infection | 1 | | | | | 1 | 0.2% |
| Teething | 1 | | | | | 1 | 0.2% |

| | | | | | | | |
|-----------------------------|------------|------------|------------|------------|-----------|------------|-------|
| Tinea | 12 | 6 | 3 | 8 | 1 | 30 | 7.4% |
| Tuberculosis | 1 | | | | | 1 | 0.2% |
| Umbilical hernia | | | | 2 | | 2 | 0.5% |
| Upper respiratory infection | 12 | 25 | 12 | 16 | 4 | 69 | 17.0% |
| Urinary tract infection | 2 | 3 | | 2 | | 7 | 1.7% |
| Vaginitis | 2 | 4 | 2 | | | 8 | 2.0% |
| Viral syndrome | 1 | | | | | 1 | 0.2% |
| Well child | 7 | 9 | 25 | 9 | 1 | 51 | 12.6% |
| Worms | 46 | 58 | 31 | 30 | 16 | 181 | 44.6% |
| TOTALS | 153 | 191 | 123 | 111 | 27 | 605 | |

***NOTE:** The percentage for each diagnosis is calculated on the basis of total patients seen in the pediatric clinics (406), not on the total number of diagnoses (because many patients had multiple diagnoses).
